# **SKYRIZI GETTING STARTED CHECKLIST**

Use this checklist from Skyrizi Complete to start and stay on track with your prescribed treatment plan.

<b>NAVIGATE INSURANCE AND SAVINGS</b> Skyrizi Complete can help you understand your insurance and find possible ways to save.	Your Nurse Ambassador is:  Your Nurse Ambassador's phone number:		
If you have not talked with your Skyrizi Complete Nurse Ambassador* yet, reach out by calling <u>1.866.SKYRIZI</u> (1.866.759.7494)			
Ask your Nurse Ambassador about your <u>savings options</u> Find out if your SKYRIZI could be as little as \$5 <sup>+</sup> per treatment			
<b>PREPARE FOR YOUR INFUSIONS</b> You'll have 3 infusions; 1 every 4 weeks. Help understanding the infusion phase of your treatment is just a click away.	Infusion location:		
Watch the SKYRIZI Infusion Video	Phone number:		
<b>Review the Infusion Checklist</b> with your Nurse Ambassador	Infusion dates: Complete		
Talk to your doctor about lab tests before, during, and up to 12 weeks of treatment with SKYRIZI	1. / / 2. / /		
Write down any questions you may have for your next doctor appointment:	3/ / [		
<b>GET READY TO INJECT AT HOME</b> You can ask for supplemental injection training to be delivered in person or during a virtual visit with your Nurse Ambassador.	Specialty Pharmacy:		
Watch the Injection Training Video			
Order resources you need for injecting at home, like the Sharps Container, Mail-back Disposal Kit, and insulated Travel Case	Phone number:		
Date of first injection at home: / /			

# Skyrizi<sup>®</sup>COMPLETE

## Not enrolled? Call 1.866.SKYRIZI (1.866.759.7494) to join today.

Once enrolled, you can expect a call from your Nurse Ambassador within 1 business day. The call may come from any area code.

\*Nurse Ambassadors are provided by AbbVie and do not work under the direction of your health care professional (HCP) or give medical advice. They are trained to direct patients to their HCP for treatment-related advice, including further referrals.

<sup>†</sup>For eligible, commercially insured patients only. See full Terms and Conditions on back. If eligible, you'll receive your Savings Card in the mail. Call your Ambassador if you do not receive your card.

Please see <u>Use and Important Safety Information</u> on page 2. Please see full <u>Prescribing Information</u>, including <u>Medication Guide</u>, at <u>https://www.rxabbvie.com/pdf/skyrizi\_pi.pdf</u> and discuss with your doctor.



# **USE AND IMPORTANT SAFETY INFORMATION** ABOUT SKYRIZI<sup>®</sup> (risankizumab-rzaa)<sup>1</sup>

# SKYRIZI USE<sup>1</sup>

SKYRIZI is a prescription medicine used to treat moderate to severe Crohn's disease in adults.

# **IMPORTANT SAFETY INFORMATION**<sup>1</sup>

### What is the most important information I should know about SKYRIZI<sup>®</sup> (risankizumab-rzaa)? SKYRIZI is a prescription medicine that may cause serious side effects, including:

# Serious allergic reactions:

- Stop using SKYRIZI and get emergency medical help right away if you get any of the following symptoms of a serious allergic reaction:
  - fainting, dizziness, feeling lightheaded (low blood pressure) - swelling of your face,
- trouble breathing or throat tightness
- chest tightness
- eyelids, lips, mouth, tongue, or throat
- skin rash, hives - itchina

### Infections:

SKYRIZI may lower the ability of your immune system to fight infections and may increase your risk of infections. Your healthcare provider should check you for infections and tuberculosis (TB) before starting treatment with SKYRIZI and may treat you for TB before you begin treatment with SKYRIZI if you have a history of TB or have active TB. Your healthcare provider should watch you closely for signs and symptoms of TB during and after treatment with SKYRIZI.

- Tell your healthcare provider right away if you have an infection or have symptoms of an infection, including:
  - fever, sweats, or chills
- weight loss

- couah

- shortness of breath
- diarrhea or stomach pain
- blood in your mucus (phlegm)
- muscle aches
- burning when you urinate or urinating more often than normal
- warm, red, or painful skin or sores on your body different from your psoriasis

Do not use SKYRIZI if you are allergic to risankizumab-rzaa or any of the ingredients in SKYRIZI.

### Before using SKYRIZI, tell your healthcare provider about all of your medical conditions, including if you:

- have any of the conditions or symptoms listed in the section "What is the most important information I should know about SKYRIZI?"
- have an infection that does not go away or that keeps coming back.
- have TB or have been in close contact with someone with TB.
- have recently received or are scheduled to receive an immunization (vaccine). Medicines that interact with the immune system may increase your risk of getting an infection after receiving live vaccines. You should avoid receiving live vaccines right before, during, or right after treatment with SKYRIZI. Tell your healthcare provider that you are taking SKYRIZI before receiving a vaccine.
- are pregnant or plan to become pregnant. It is not known if SKYRIZI can harm your unborn baby.

Please see full Prescribing Information, including Medication Guide,

at https://www.rxabbvie.com/pdf/skyrizi\_pi.pdf and discuss with your doctor.

 are breastfeeding or plan to breastfeed. It is not known if SKYRIZI passes into your breast milk.

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements.

## What are the possible side effects of SKYRIZI?

### SKYRIZI may cause serious side effects. See "What is the most important information I should know about SKYRIZI?"

Liver problems in Crohn's disease: A person with Crohn's disease who received SKYRIZI by intravenous infusion developed changes in liver blood tests with a rash that led to hospitalization. Your doctor will do blood tests to check your liver before, during, and up to 12 weeks of treatment and may stop treatment with SKYRIZI if you develop liver problems. Tell your doctor right away if you notice any of the following symptoms: unexplained rash, nausea, vomiting, stomach (abdominal) pain, tiredness (fatigue), loss of appetite, yellowing of the skin and eyes (jaundice), and dark urine.

The most common side effects of SKYRIZI in people treated for Crohn's disease include: upper respiratory infections, injection site reactions. fever, headache, stomach (abdominal) pain, back pain, joint pain, and low red blood cells (anemia).

These are not all the possible side effects of SKYRIZI. Call your doctor for medical advice about side effects.

Use SKYRIZI exactly as your healthcare provider tells you to use it.

SKYRIZI is available in a 600 mg/10 mL intravenous infusion and a 360 mg/2.4 mL single-dose prefilled cartridge with on-body injector.

### You are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch or call 1-800-FDA-1088.

If you are having difficulty paying for your medicine, AbbVie may be able to help. Visit AbbVie.com/myAbbVieAssist to learn more.

Reference: 1. SKYRIZI [package insert]. North Chicago, IL: AbbVie Inc.

# **SKYRIZI COMPLETE SAVINGS CARD TERMS & CONDITIONS**

Eligibility: Available to patients with commercial insurance coverage for SKYRIZI<sup>®</sup> (risankizumab-rzaa) who meet eligibility criteria. This co-pay assistance program is not available to patients receiving prescription reimbursement under any federal, state, or government-funded insurance programs (for example, Medicare [including Part D], Medicare Advantage, Medigap, Medicaid, TRICARE, Department of Defense, or Veterans Affairs programs) or where prohibited by law. Offer subject to change or termination without notice. Restrictions, including monthly maximums, may apply. This is not health insurance. For full Terms and Conditions, visit SKYRIZICDSavingsCard.com. To learn about AbbVie's privacy practices and your privacy choices, visit www.abbvie.com/privacy.html



© 2022 AbbVie Inc. All rights reserved. SKYRIZI® and its design are registered trademarks of AbbVie Biotechnology Ltd. US-SKZ-210530 June 2022

# Skyrizi<sup>® COMPLETE</sup>

SCD-032022-AP02

## Fax to Skyrizi Complete (1.678.727.0690) Questions? Call 1.866.759.7494

# **Enrollment and Prescription Form**

Sections in BLUE (1, 2, 3, 4) are necessary for enrollment into Skyrizi Complete. Required fields are marked with an asterisk (\*).

**1** PATIENT'S INFORMATION - The health care professional (HCP) and the patient should fill out this form completely before leaving the office.

Please print clearly.							
First Name*:	Last Name*:	C	Date of Birth:	/ /	Gen	der: (check o	one) 🗆 M 🗆 F
Address*:		City*:			Stat	e*: Z	ZIP*:
Home Phone*:	Mobile Phone:	Email Address*:			□Sp	anish interp	reter needed
mobile number. Message a	ated and recurring text messages from Al and data rates may apply. My consent is n v full Terms and Conditions.						
BestTime to Call (Monday-Fi	iday): 🗌 Anytime 🗌 Morning 🗌 A	fternoon 🗌 Evening					
What was patient's last compl	eted treatment? 🗌 Not started 🔲 Infusior	1 $\Box$ Infusion 2 $\Box$ Infusion 3	SKYRIZI On-B	ody Injector	Date of Last Tre	atment:	//
professional (HCP) or give med collected on this form include about the categories of perso	Your own Support Specialist/Nurse provided ical advice. They are trained to direct patient a name, date of birth, address, Rx, and ins inal information collected by AbbVie and s and updates about AbbVie's products, c	s to their HCP for treatment-relat surance information. The inform the purposes for which AbbVie	ed advice, includi nation collected v uses personal in	ng further ref vill be used oformation, v	errals. The categ for program enro visit <u>https://priv</u>	ories of pers ollment. For <u>acy.abbvie</u>	sonal information more information
INSURANCE INFORMAT	ON □ Check box if you will attach copy	of Insurance Cards. Please als	so provide suppl	emental insi	urance.		
Demoficien /Condhelder Nore		: Desseriation In					

Beneficiary/Cardholder Name:	Prescription Insurance:						
Medical Insurance:	Rx Group #:	Rx Group #:					
Medical Insurance ID #:							
Group #:	Rx Bin #:	Rx PCN #:					
	▼ FOR HEALTH CARE PROVIDER USE ONLY ▼						
3 DIAGNOSIS* Crohn's disease (CD) ICD-10:	Date of Diagnosis://						
	eive a copy:	3					
NPI #*:	Office Contact Name:						
	Office Fax*:	_ :					
5 CLINICAL INFORMATION Prior Therapies:	Concomitant Medications: Drug Allergies:	TBTest (Date):/					
·	Drug Allergies.	_ Specialty Pharmacy only.					
6 SITE OF INFUSION INFORMATION							
Prescriber's office (if checked, skip to section 7) Physician or Infusion Provider Name:	Address:	Infusion center Other:					
Practice/Facility Name:		State: ZIP: Fax:					
8 PRESCRIPTION INFORMATION - Fill out and sig	cy prescription 8a below. gn corresponding prescription(s) below						
Prescriptions will be forwarded to pharmacy	8a. PHARMACY PRESCRIPTION - OPTIONAL						
designated in section 8a or to a pharmacy within the patient's approved network, if blank.	Patient's preferred Specialty Pharmacy:						
Pharmacy may call to verify the prescription.	PRESCRIBER CERTIFICATION: I certify that the above therapy is medically necessary and that the information provided is accurate to the best of my knowledge. I certify that I am the prescriber who has prescribed SKYRIZI to the previously identified patient and that I provided the patient with a description of the Skyrizi Complete patient support program. I authorize Skyrizi Complete to act on my behalf for the purposes of						
Check appropriate boxes to indicate quantity to dispense and directions:	transmitting this prescription to the appropriate pharmacy designated	by the patient utilizing their benefit plan (if applicable).					
	Prescriber's Signature: (REQUIRED)	Date://					
Initiation Therapy <sup>+</sup> —SKYRIZI 600 mg/10 mL single use vial Week 0: 600 mg to be administered via IV Infusion 1 vial; no refills	8b. SKYRIZI COMPLETE PRESCRIPTION - required in the event a commercially insured patient with a valid Rx for SKYRIZI experiences an insurance delay or denial						
Uweek 4: 600 mg to be administered via IV Infusion 1 vial; no refills	Prescription to be filled through an AbbVie-authorized pharmacy. I unde						
<ul> <li>Week 8: 600 mg to be administered via IV Infusion 1 vial; no refills</li> <li><sup>1</sup>For the treatment of Crohn's disease, evaluate liver enzymes and bilirubin at baseline, and during induction at least up to 12 weeks of treatment. Monitor thereafter according to routine patient management.</li> </ul>	PRESCRIBER CERTIFICATION: I certify that I am the prescriber who has prescribed SKYRIZI to the previously identified patient and that I provided the patient with a description of the Skyrizi Complete patient support program. I authorize Skyrizi Complete to act on my behalf for the purposes of transmitting this prescription to the appropriate pharmacy. I understand that the no charge resource through Skyrizi Complete may support patients who are experiencing a delay in insurance coverage for SKYRIZI until coverage is obtained, and I confirm that I will support the above-identified patient in seeking to secure such coverage as I deem appropriate. I certify that I will not seek reimbursement from any third party payor for any no charge product dispensed by an AbbVie-authorized pharmacy. I confirm my patient has or will complete IV initiation therapy as recommended in the FDA-approved prescribing information.						
Ongoing Therapy—SKYRIZI 360 mg/2.4 mL prefilled cartridge with SKYRIZI On-Body Injector	initiation therapy as recommended in the FDA-approved prescribing	information.					
Week 12: Inject 360 mg SC and every 8 weeks thereafter 1 device with prefilled cartridge; Refills:	Prescriber's Signature: (REQUIRED)	Date:/					

IMPORTANT INFORMATION: By submitting this form, you are referring the above patient to AbbVie's patient support program to determine eligibility and receive support related to an AbbVie product. The personal information collected on this form will be used for program management and to perform research and analytics. For more information about the categories of personal information collected by AbbVie and the purposes for which AbbVie uses personal information, visit <u>https://privacy.abbvie</u>. Please share this information with your patient.



US-SKZ-210530

# Indication and Important Safety Information<sup>1</sup>

# SKYRIZI INDICATION<sup>1</sup>

SKYRIZI is indicated for the treatment of moderately to severely active Crohn's disease in adults.

## **IMPORTANT SAFETY INFORMATION**<sup>1</sup>

### **Hypersensitivity Reactions**

SKYRIZI<sup>®</sup> (risankizumab-rzaa) is contraindicated in patients with a history of serious hypersensitivity reaction to risankizumab-rzaa or any of the excipients. Serious hypersensitivity reactions, including anaphylaxis, have been reported with the use of SKYRIZI. If a serious hypersensitivity reaction occurs, discontinue SKYRIZI and initiate appropriate therapy immediately.

### Infection

SKYRIZI may increase the risk of infection. Do not initiate treatment with SKYRIZI in patients with a clinically important active infection until it resolves or is adequately treated.

In patients with a chronic infection or a history of recurrent infection, consider the risks and benefits prior to prescribing SKYRIZI. Instruct patients to seek medical advice if signs or symptoms of clinically important infection occur. If a patient develops such an infection or is not responding to standard therapy, closely monitor and discontinue SKYRIZI until the infection resolves.

### **Tuberculosis (TB)**

Prior to initiating treatment with SKYRIZI, evaluate for TB infection and consider treatment in patients with latent or active TB for whom an adequate course of treatment cannot be confirmed. Monitor patients for signs and symptoms of active TB during and after SKYRIZI treatment. Do not administer SKYRIZI to patients with active TB.

### Hepatotoxicity in Treatment of Crohn's Disease

Drug-induced liver injury was reported in a patient with Crohn's disease who was hospitalized for a rash during induction dosing of SKYRIZI. For the treatment of Crohn's disease, evaluate liver enzymes and bilirubin at baseline and during induction (12 weeks); monitor thereafter according to routine patient management. Consider an alternate treatment for patients with evidence of liver cirrhosis. Interrupt treatment if drug-induced liver injury is suspected, until this diagnosis is excluded. Instruct your patient to seek immediate medical attention if they experience symptoms suggestive of hepatic dysfunction.

### **Administration of Vaccines**

Avoid use of live vaccines in patients treated with SKYRIZI. Medications that interact with the immune system may increase the risk of infection following administration of live vaccines. Prior to initiating SKYRIZI, complete all age appropriate vaccinations according to current immunization guidelines.

### **Adverse Reactions**

Most common (>3%) adverse reactions associated with SKYRIZI in Crohn's disease are upper respiratory infections, headache, and arthralgia in induction and arthralgia, injection site reactions, abdominal pain, anemia, pyrexia, back pain, arthropathy, and urinary tract infection in maintenance.

**Lipid Elevations**: Increases from baseline and increases relative to placebo were observed at Week 4 and remained stable to Week 12 in patients treated with SKYRIZI in Crohn's disease.

**Dosage Forms and Strengths**: SKYRIZI is available in a 600 mg/10 mL intravenous infusion and a 360 mg/2.4 mL single-dose prefilled cartridge with on-body injector.

Reference: 1. SKYRIZI [package insert]. North Chicago, IL: AbbVie Inc.

## **Skyrizi Complete Prescription Terms & Conditions**

**Eligibility criteria:** Available to patients aged 63 or younger with commercial insurance coverage. Patients must have a valid prescription for SKYRIZI<sup>®</sup> (risankizumab-rzaa) for an FDA-approved indication, and a denial of insurance coverage based on a prior authorization request on file, and a confirmation of appeal. For medical coverage, a pre-certification submission will be required. Continued eligibility for the program requires the submission of an appeal of the coverage denial every 180 days. Program provides for SKYRIZI<sup>®</sup> (risankizumab-rzaa) at no charge to patients for up to two years or until they receive insurance coverage approval, whichever occurs earlier, and is not contingent on purchase requirements of any kind. Program is not available to patients whose medications are reimbursed in whole or in part by Medicare, Medicaid, TRICARE, or any other federal or state program. Offer subject to change or discontinuance without notice. This is not health insurance, and program does not guarantee insurance coverage. No claims for payment may be submitted to any third party for product dispensed by program. Limitations may apply.

Skyrizi risankizumab-rzaa

Please see full Prescribing Information.

